



CORIELL INSTITUTE

FOR MEDICAL RESEARCH

GM23392*B

Certificate of Analysis

Product description	Human Fibroblast reprogrammed with four factors(OCT4, SOX2, NANOG, LIN28A) using retroviral vector	
Publication(s) describing iPSC establishment		
Parent cell line and cell type	GM06114	Fibroblast
Diagnosis	Apparently Healthy Fetal Tissue	
Passage of iPSC reported at submission	16	
Number of passages at Coriell	5	
Media	DMEM/F12 + 20% KOSR + 100ng/ml FGF	
Feeder	CF1 MEFs on 0.1% Gelatin	
Passage method	Collagenase	
Split ratio	1:6; every 5-7 days	

The following testing specifications have been met for the specified product lot:

Test Description	Test Method	Test Specification	Result
Post-Thaw Viable Cell Recovery	Colony Doubling	Colony formation and diameter doubling within 5 days	Pass
Sterility	Growth on agar	Negative	Pass
Mycoplasma	PCR	Negative	Pass
Karyotype	G-banding	46 XY	Pass
Identity Match	STR (THO-1, D22S417, D10S526, vWA31, D5S592, and FES/FPS)	Match parent fibroblast line	Pass
Surface Antigen Expression of Stem Cell Markers	Immunostaining	> 80% expression of SSEA-4	Pass
Teratoma Formation	<i>In Vivo</i> Teratoma formation	3 germ layer teratoma	Pass

Post-Thaw Viability

One vial of distribution lot was thawed. Cultures were observed daily. Colonies were photographed when they first appeared, then 7 days later (Colonies must double in diameter within 5 days).

Day 2	222 μm
Day 9	1069 μm




Figure 1A. Colony observed post thaw




Figure 1B. Colony 7 days after first observation

Karyotype Analysis




Figure 2: G-banded karyotype showing 46 XY

Surface Antigen Expression of Stem Cell Markers

Undifferentiated cells are stained for the surface antigen, SSEA4. SSEA4 (stage specific embryonic antigen 4) is expressed on undifferentiated human stem cells.




Figure 3: Dot Plot of SSEA-4 positive population.

Assessment of Pluripotency of a Cell Line

Cells are directed to differentiate to assess the pluripotency of the cell line. RNA is harvested and gene expression is analyzed by real-time PCR. Ct values are adjusted for loading using a housekeeping gene. Gene expression is shown as fold difference to undifferentiated cells.

Embryoid Body (EB) Formation Assay




Figure 4. Gene expression following EB differentiation. Fold difference is shown relative to undifferentiated iPS cell line.

Pluripotency Markers

	OCT4	SOX2	NANOG	GDF3	REXO1
GM23392*B EB	-14	-3	-11	-5	-1

Ectoderm

	PAX6	NES	TP63	KRT14	NOG
GM23392*B EB	85	-1	283	294	12

Mesoderm

	T	RUNX1	DES	PECAM1	TAL1
GM23392*B	-1	11	-5	3	-2

Endoderm

	SOX7	SOX17	FOXA2	AFP
GM23392*B	46	-6	-6	3583

Table 1. Fold difference values of gene expression of EB. Fold difference is relative to undifferentiated cells. Ct values are normalized to that of GAPDH.

Neural Differentiation




Figure 5A. Image of Neuronal Differentiation




Figure 5B. Gene expression following neuronal differentiation. Fold difference is shown relative to undifferentiated iPS cell line.

Cardiac Differentiation




Figure 6A. Image of cardiac differentiation.




Figure 6B. Gene expression following cardiac differentiation. Fold difference is shown relative to undifferentiated iPS cell line.

Definitive Endoderm Differentiation




Figure 7A. Image of Definitive Endoderm Differentiation





Figure 7B. Gene expression following Definitive Endoderm differentiation. Fold difference is shown relative to undifferentiated iPS cell line.

- Pass
- Fail
- Other:



Steve Madore, PhD
Director, Stem Cell Biobank
July 27, 2012

Teratoma Formation Analysis Report

Project Information

Service title: Teratoma Formation Analysis
 Customer: Coriell Institute
 PI/Contact person: Karen Fecenko-Tacka
 Report date: July 17, 2012
 Project manager: Qi Zheng
 Contact person: Tianmin "Ivy" Zhang

Service Detail

Cell type: human iPS cells
 Cell line & passage: GM23392, P7
 Feeder layer: CF1 MEF
 Mouse type: Fox Chase SCID-beige, male, 6 week old, from Charles River
 Cell concentration: 2-3 million/site, in 30% Matrigel
 3 H&E slides
 Injection date: May 10, 2012

	Mouse #1	Mouse #2	Mouse #3	Control
Injection sites	kidney capsule testis	kidney capsule testis	kidney capsule testis	kidney capsule testis
Tissue harvested	one kidney tumor and one testis tumor	one kidney tumor and one testis tumor	one kidney tumor and one testis tumor	one kidney tumor and one testis tumor
Days post-injection	60	60	60	60


H&E Histology Instruction

Histology: 10% Formalin fixed over night, embedded in paraffin, cut into 5- μ m serial sections, H&E staining
 Three embryonic germ cell layers: endoderm, mesoderm and ectoderm
 Endoderm: digestive system (includes liver and pancreas), respiratory system, most glands
 Mesoderm: muscle, blood vessels, much of the genital-urinary system, skeletal system
 Ectoderm: skin, hair, nails, sweat and mammary glands, nervous system (including hypothalamus and both lobes of the pituitary gland), oral and nasal cavities, portions of the vagina, vestibule, penis and clitoris

Tumor and organ pictures



Mouse#1: one kidney tumor (left) and one testis tumor (right) harvested on day 60 after injection




Mouse#2: one kidney tumor (left) and one testis tumor (right) harvested on day 60 after injection



Mouse#3: one kidney tumor(left) and one testis tumor (right) harvested on day 60 after injection


H&E staining results of kidney and testis tumors:

Endoderm




Gland (200x)


Mesoderm




Cartilage (200x)




Bone (200x)



Adipose tissue (200x)




Muscle (200x)




Blood vessel (200x)

Ectoderm



Neuronal rosette (200x)



Pigmented cells (200x)

Summary

Three kidney tumors and three testis tumors are composed of scattered regions of differentiated cells and a large population of undifferentiated neoplastic cells. Three germ layers were clearly identified in histology analysis. The tissues listed above indicate that small areas of what might be these kinds of tissues were noted within the tumor. Overall, there is some degree of differentiation of these cells with organized structures, suggesting that some of these cells are pluripotent.

Project manager

Signature: Date: 7/17/2012Qi Zheng, Ph.D.
Senior Scientist

Reviewed and proved by

Signature: Date: 7/17/2012Steve Yu, Ph.D.
Director of Service Department